

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 186 672 B1**

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
**30.11.2005 Bulletin 2005/48**

(51) Int Cl.<sup>7</sup>: **C12Q 1/68**, C07K 14/47,  
C07K 16/18

(21) Application number: **01306983.6**

(22) Date of filing: **17.08.2001**

### (54) Polymorphisms in the human organic anion transporter C (OATP-C) gene

Polymorphismen im Gen für den humanen organischen Anionentransporter C (OATP-C)

Polymorphismes dans le gène humain du transporteur d'anion organique (OATP-C)

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR**

(30) Priority: **23.08.2000 US 226909 P**

(43) Date of publication of application:  
**13.03.2002 Bulletin 2002/11**

(60) Divisional application:  
**05011183.0 / 1 589 115**

(73) Proprietor: **AstraZeneca AB**  
**151 85 Södertälje (SE)**

(72) Inventors:  
• **Adeokun, Monisola**  
Cheshire, SK10 4TG (GB)  
• **Ambrose, Helen Jean**  
Cheshire, SK10 4TG (GB)  
• **Cresswell, Carl John**  
Cheshire, SK10 4TG (GB)  
• **Dudley, Adam Jeston**  
Wilmington, DE 19850-5437 (US)

(74) Representative: **Giles, Allen Frank et al**  
**AstraZeneca PLC**  
Global Intellectual Property  
Mereside  
Alderley Park  
Macclesfield, Cheshire SK10 4TG (GB)

(56) References cited:  
**WO-A-00/08157 WO-A-99/06046**

- **TAMAI I ET AL: "MOLECULAR IDENTIFICATION AND CHARACTERIZATION OF NOVEL MEMBERS OF THE HUMAN ORGANIC ANION TRANSPORTER (OATP) FAMILY" BIOCHIMICA ET BIOPHYSICA ACTA, AMSTERDAM, NL, vol. 273, no. 1, 2000, pages 251-260, XP000941538 ISSN: 0006-3002 & DATABASE EMBL [Online] EBI; 8 June 1999 (1999-06-08) TAMAI I ET AL.: "Homo sapiens mRNA for organic anion transporter OATP-C" retrieved from [HTTP://WWW.EBI.AC.UK/CGI-BIN/EMBLFETCH](http://www.ebi.ac.uk/cgi-bin/emblfetch) Database accession no. AB026257**
- **DATABASE EMBL [Online] EBI; 2 February 2000 (2000-02-02) MUZNY DM ET AL.: "Homo sapiens 12p BAC RP11-12505" retrieved from [HTTP://WWW.EBI.AC.UK/CGI-BIN/EMBLFETCH](http://www.ebi.ac.uk/cgi-bin/emblfetch) Database accession no. AC022335 XP002188259**
- **HSIANG B ET AL: "A NOVEL HUMAN HEPATIC ORGANIC ANION TRANSPORTING POLYPEPTIDE (OATP2)" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 274, no. 52, 1999, pages 37161-37168, XP000877290 ISSN: 0021-9258**
- **ULBRECHT M ET AL.: "Association of beta 2-adrenoreceptor variants with bronchial hyperresponsiveness" AMERICAN JOURNAL OF RESPIRATORY AND CRITICAL CARE MEDICINE, vol. 161, no. 2, February 2000 (2000-02), pages 469-474, XP002188594**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 1 186 672 B1**

## Description

**[0001]** This invention relates to polymorphisms in the human OATPC gene and corresponding novel allelic polypeptides encoded thereby. The invention also relates to methods and materials for analysing allelic variation in the OATPC gene, and to the use of OATPC polymorphism in treatment of diseases with OATPC transportable drugs.

**[0002]** Na<sup>+</sup>-independent organic anion transporting polypeptide (OATP) C gene is a member of the OATP supergene family involved in multifunctional transport of organic anion. OATPC transports the organic anion taurocholate, conjugated steroids: DHEAS, estradiol 17 $\beta$ -D-glucuronide and estrone-3-sulfate, eicosanoids: PGE<sub>2</sub>, thromboxane B<sub>2</sub>, leukotriene C<sub>4</sub>, and E<sub>4</sub>, and thyroid hormones T<sub>4</sub> and T<sub>3</sub><sup>1,2</sup>. OATPC has also been shown to be involved in the transport of xenobiotics, and drugs involved in lipid lowering *e.g.* statins<sup>1</sup>. Statins have been referred to as a first-line therapy for patients with atherosclerotic vascular diseases. The OATPC gene and its product is also thought to be of importance in other diseases due to its transport of DHEAS an adrenal steroid which has been suggested to have positive neuropsychiatric, immune, and metabolic effects<sup>3</sup>. Due to the substrate specificity, location in the liver, and being exclusively expressed in the liver, Abe *et al* suggested that OATPC could be the predominant clearance mechanism of several endogenous and exogenous substrates in the liver. OATPC is the first human molecule reported to transport thyroid hormones<sup>2</sup>.

**[0003]** This liver specific transporter may be useful in liver-specific drug delivery systems and liver-specific chemotherapy, bile acid formation and the pathogenesis of diseases such as cholestasis, hyperbilirubinemia and thyroid hormone resistance.

**[0004]** The OATPC gene (sometimes called OAPT2 in the literature) has been cloned by four different groups, annotated and published as EMBL accession numbers AB026257 (OATPC, 2452bp), AF205071 (OATP2, 2830, ref 1), AJ132573 (OATP2, 2778)<sup>4</sup> and AF060500 (LST-1)<sup>2</sup>. Polymorphism has been reported by Tamai<sup>5</sup> which is Asn 130Asp and Val174Ala although any functional effect was stated therein to be not clear. König (2000) J Biol Chem 275, 23161-23168 describes the genomic organisation of OATP 1, 2 and 8. International patent application WO 00/08157 describes human anion transporter genes and some polymorphisms.

**[0005]** All positions herein of polymorphisms in the OATPC polynucleotide relate to the position in one of SEQ ID NO 1 or 3-12 unless stated otherwise or apparent from the context.

**[0006]** All positions herein of polymorphisms in the OATPC polypeptide relate to the position in SEQ ID NO 2 unless stated otherwise or apparent from the context.

**[0007]** One approach is to use knowledge of polymorphisms to help identify patients most suited to therapy with particular pharmaceutical agents (this is often termed "pharmacogenetics"). Pharmacogenetics can also be used in pharmaceutical research to assist the drug selection process. Polymorphisms are used in mapping the human genome and to elucidate the genetic component of diseases. The reader is directed to the following references for background details on pharmacogenetics and other uses of polymorphism detection: Linder *et al.* (1997), Clinical Chemistry, **43**, 254; Marshall (1997), Nature Biotechnology, **15**, 1249; International Patent Application WO 97/40462, Spectra Biomedical; and Schafer *et al.* (1998), Nature Biotechnology, **16**, 33.

**[0008]** Clinical trials have shown that patient response to treatment with pharmaceuticals is often heterogeneous. Thus there is a need for improved approaches to pharmaceutical agent design and therapy.

**[0009]** Point mutations in polypeptides will be referred to as follows: natural amino acid (using 1 or 3 letter nomenclature), position, new amino acid. For (a hypothetical) example "D25K" or "Asp25Lys" means that at position 25 an aspartic acid (D) has been changed to lysine (K). Multiple mutations in one polypeptide will be shown between square brackets with individual mutations separated by commas.

**[0010]** The present invention is based on the discovery of polymorphisms in OATPC.

**[0011]** According to one aspect of the present invention there is provided a method for the detection of a polymorphism in OATPC in a human, which method comprises determining the sequence of the human at:

position 1561 of the OATPC gene as defined by the position in SEQ ID NO: 1 is not G; or  
position 488 in OATPC polypeptide defined by position in SEQ ID NO: 2 is not Gly.

**[0012]** The term human includes both a human having or suspected of having a OATPC mediated disease and an asymptomatic human who may be tested for predisposition or susceptibility to such disease. At each position the human may be homozygous for an allele or the human may be a heterozygote.

<sup>2</sup> Identification of a Novel Gene Family Encoding Human liver-specific Organic Anion Transporter LST-1, Takaaki Abe *et al* J Biol Chem **274**, 17159-17163 (1999)

<sup>1</sup> A Novel Human Hepatic Organic Anion Transporting Polypeptide (OATP2), Hsiang *et al* J Biol Chem **274**, 37161-37168 (1999)

<sup>3</sup> Bates *et al* (1998) Curr. Opin. Endocrinol. Diab. **5**, 357-366

<sup>4</sup> A novel human organic anion transporting polypeptide localised to the basolateral hepatocyte membrane, König Jorg *et al* (2000) Am J Physiol. Gastrointest. Liver Physiol. **278**: G156-G164

<sup>5</sup> Tamai *et al* (2000), BBRC, 273, 251-60

[0013] The term polymorphism includes single nucleotide substitution, nucleotide insertion and nucleotide deletion which in the case of insertion and deletion includes insertion or deletion of one or more nucleotides at a position of a gene.

[0014] The method for diagnosis is preferably one in which the sequence is determined by a method selected from amplification refractory mutation system and restriction fragment length polymorphism.

[0015] The status of the individual may be determined by reference to allelic variation at any one, two, three, four, five, six, seven, eight, nine or more positions.

[0016] The test sample of nucleic acid is conveniently a sample of blood, bronchoalveolar lavage fluid, sputum, or other body fluid or tissue obtained from an individual. It will be appreciated that the test sample may equally be a nucleic acid sequence corresponding to the sequence in the test sample, that is to say that all or a part of the region in the sample nucleic acid may firstly be amplified using any convenient technique e.g. PCR, before analysis of allelic variation.

[0017] It will be apparent to the person skilled in the art that there are a large number of analytical procedures which may be used to detect the presence or absence of variant nucleotides at one or more polymorphic positions of the invention. In general, the detection of allelic variation requires a mutation discrimination technique, optionally an amplification reaction and optionally a signal generation system. Table 1 lists a number of mutation detection techniques, some based on the PCR. These may be used in combination with a number of signal generation systems, a selection of which is listed in Table 2. Further amplification techniques are listed in Table 3. Many current methods for the detection of allelic variation are reviewed by Nollau *et al.*, Clin. Chem. **43**, 1114-1120, 1997; and in standard textbooks, for example "Laboratory Protocols for Mutation Detection", Ed. by U. Landegren, Oxford University Press, 1996 and "PCR", 2<sup>nd</sup> Edition by Newton & Graham, BIOS Scientific Publishers Limited, 1997.

#### Abbreviations:

[0018]

ALEX™	Amplification refractory mutation system linear extension
APEX	Arrayed primer extension
ARMS™	Amplification refractory mutation system
b-DNA	Branched DNA
bp	base pair
CMC	Chemical mismatch cleavage
COPS	Competitive oligonucleotide priming system
DGGE	Denaturing gradient gel electrophoresis
FRET	Fluorescence resonance energy transfer
HMG-CoA	3-hydroxy-3-methylglutaryl-coenzyme A
LCR	Ligase chain reaction
MASDA	Multiple allele specific diagnostic assay
NASBA	Nucleic acid sequence based amplification
OATP	Na <sup>+</sup> -independent organic anion transporting polypeptide
OLA	Oligonucleotide ligation assay
PCR	Polymerase chain reaction
PTT	Protein truncation test
RFLP	Restriction fragment length polymorphism
SDA	Strand displacement amplification
SNP	Single nucleotide polymorphism
SSCP	Single-strand conformation polymorphism analysis
SSR	Self sustained replication

(continued)

TGGE	Temperature gradient gel electrophoresis
------	--

Table 1 - Mutation Detection Techniques

**[0019] General:** DNA sequencing, Sequencing by hybridisation

**Scanning:** PTT\*, SSCP, DGGE, TGGE, Cleavase, Heteroduplex analysis, CMC, Enzymatic mismatch cleavage

\* Note: not useful for detection of promoter polymorphisms.

#### Hybridisation Based

**[0020]** Solid phase hybridisation: Dot blots, MASDA, Reverse dot blots, Oligonucleotide arrays (DNA Chips)

Solution phase hybridisation: Taqman™ - US-5210015 & US-5487972 (Hoffmann-La Roche), Molecular Beacons - Tyagi *et al* (1996), Nature Biotechnology, 14, 303; WO 95/13399 (Public Health Inst., New York)

**Extension Based:** ARMS™, ALEX™ - European Patent No. EP 332435 B1 (Zeneca Limited), COPS - Gibbs *et al* (1989), Nucleic Acids Research, 17, 2347.

**Incorporation Based:** Mini-sequencing, APEX

**Restriction Enzyme Based:** RFLP, Restriction site generating PCR

**Ligation Based:** OLA

**Other:** Invader assay

Table 2 - Signal Generation or Detection Systems

**[0021] Fluorescence:** FRET, Fluorescence quenching, Fluorescence polarisation - United Kingdom Patent No. 2228998 (Zeneca Limited)

**Other:** Chemiluminescence, Electrochemiluminescence, Raman, Radioactivity, Colorimetric, Hybridisation protection assay, Mass spectrometry

Table 3 - Further Amplification Methods

**[0022]** SSR, NASBA, LCR, SDA, b-DNA

**[0023]** Preferred mutation detection techniques include ARMS™, ALEX™, COPS, Taqman, Molecular Beacons, RFLP, and restriction site based PCR and FRET techniques.

**[0024]** Particularly preferred methods include ARMS™ and RFLP based methods. ARMS™ is an especially preferred method.

**[0025]** In a further aspect, the diagnostic methods of the invention are used to assess the pharmacogenetics of a drug transportable by OATPC.

**[0026]** Assays, for example reporter-based assays, may be devised to detect whether one or more of the above polymorphisms affect transcription levels and/or message stability.

**[0027]** Individuals who carry particular allelic variants of the OATPC gene may therefore exhibit differences in their ability to regulate protein biosynthesis under different physiological conditions and will display altered abilities to react to different diseases. In addition, differences arising as a result of allelic variation may have a direct effect on the response of an individual to drug therapy. The diagnostic methods of the invention may be useful both to predict the clinical response to such agents and to determine therapeutic dose.

**[0028]** In a further aspect, the diagnostic methods of the invention, are used to assess the predisposition and/or susceptibility of an individual to diseases mediated by OATPC. This may be particularly relevant in the development of hyperlipoproteinemia and cardiovascular disease and the present invention may be used to recognise individuals who are particularly at risk from developing these conditions.

**[0029]** In a further aspect, the diagnostic methods of the invention are used in the development of new drug therapies which selectively target one or more allelic variants of the OATPC gene. Identification of a link between a particular allelic variant and predisposition to disease development or response to drug therapy may have a significant impact on the design of new drugs. Drugs may be designed to regulate the biological activity of variants implicated in the disease process whilst minimising effects on other variants.

**[0030]** In a further diagnostic aspect of the invention the presence or absence of variant nucleotides is detected by reference to the loss or gain of, optionally engineered, sites recognised by restriction enzymes.

**[0031]** According to another aspect of the present invention there is provided a human OATPC gene or its complementary strand comprising a variant allelic polymorphism at one or more of positions defined herein or a fragment thereof of at least 20 bases comprising at least one novel polymorphism.

**[0032]** Fragments are at least 17 bases, more preferably at least 20 bases, more preferably at least 30 bases.

**[0033]** According to another aspect of the present invention there is provided a polynucleotide comprising at least 20 bases of the human OATPC gene and comprising an allelic variant selected from any one of the following:

Region	variant	Position in SEQ ID NO	SEQ ID NO
Exon 10	C	1561	1

**[0034]** According to another aspect of the present invention there is provided a human OATPC gene or its complementary strand comprising a polymorphism, preferably corresponding with one or more the positions defined herein or a fragment thereof of at least 20 bases comprising at least one polymorphism.

**[0035]** Fragments are at least 17 bases, more preferably at least 20 bases, more preferably at least 30 bases.

**[0036]** The invention further provides a nucleotide primer which can detect a polymorphism of the invention.

**[0037]** According to another aspect of the present invention there is provided an ARMS allele specific primer or an allele-specific oligonucleotide probe selected from one of the following:

an ARMS allele specific primer;  
or an allele-specific oligonucleotide probe;  
capable of detecting a OATPC gene polymorphism, preferably at one or more of the positions as defined herein.

**[0038]** An allele specific primer is used, generally together with a constant primer, in an amplification reaction such as a PCR reaction, which provides the discrimination between alleles through selective amplification of one allele at a particular sequence position e.g. as used for ARMS™ assays. The allele specific primer is preferably 17- 50 nucleotides, more preferably about 17-35 nucleotides, more preferably about 17-30 nucleotides.

**[0039]** An allele specific primer preferably corresponds exactly with the allele to be detected but derivatives thereof are also contemplated wherein about 6-8 of the nucleotides at the 3' terminus correspond with the allele to be detected and wherein up to 10, such as up to 8, 6, 4, 2, or 1 of the remaining nucleotides may be varied without significantly affecting the properties of the primer.

**[0040]** Primers may be manufactured using any convenient method of synthesis. Examples of such methods may be found in standard textbooks, for example "Protocols for Oligonucleotides and Analogues; Synthesis and Properties," Methods in Molecular Biology Series; Volume 20; Ed. Sudhir Agrawal, Humana ISBN: 0-89603-247-7; 1993; 1<sup>st</sup> Edition. If required the primer(s) may be labelled to facilitate detection.

**[0041]** According to another aspect of the present invention there is provided an allele-specific oligonucleotide probe capable of detecting a OATPC gene polymorphism, preferably at one or more of the positions defined herein.

**[0042]** The allele-specific oligonucleotide probe is preferably 17- 50 nucleotides, more preferably about 17-35 nucleotides.

**[0043]** The design of such probes will be apparent to the molecular biologist of ordinary skill. Such probes are of any convenient length such as up to 50 bases, up to 40 bases, more conveniently up to 30 bases in length, such as for example 8-25 or 8-15 bases in length. In general such probes will comprise base sequences entirely complementary to the corresponding wild type or variant locus in the gene. However, if required one or more mismatches may be introduced, provided that the discriminatory power of the oligonucleotide probe is not unduly affected. The probes of the invention may carry one or more labels to facilitate detection.

**[0044]** According to another aspect of the present invention there is provided an allele specific primer or an allele specific oligonucleotide probe capable of detecting a OATPC gene polymorphism at one of the positions defined herein.

**[0045]** In another aspect of the invention, the single nucleotide polymorphisms of this invention may be used as genetic markers in linkage studies. This particularly applies to the polymorphisms of relatively high frequency. The OATPC gene is on chromosome 12p (as shown from a database search with the cDNA as a query sequence). Low frequency polymorphisms may be particularly useful for haplotyping as described below. A haplotype is a set of alleles found at linked polymorphic sites (such as within a gene) on a single (paternal or maternal) chromosome. If recombination within the gene is random, there may be as many as 2<sup>n</sup> haplotypes, where 2 is the number of alleles at each SNP and n is the number of SNPs. One approach to identifying mutations or polymorphisms which are correlated with clinical response is to carry out an association study using all the haplotypes that can be identified in the population of interest. The frequency of each haplotype is limited by the frequency of its rarest allele, so that SNPs with low frequency alleles are particularly useful as markers of low frequency haplotypes. As particular mutations or polymorphisms associated with certain clinical features, such as adverse or abnormal events, are likely to be of low frequency

within the population, low frequency SNPs may be particularly useful in identifying these mutations (for examples see: Linkage disequilibrium at the cystathionine beta synthase (CBS) locus and the association between genetic variation at the CBS locus and plasma levels of homocysteine. *Ann Hum Genet* (1998) **62**:481-90, De Stefano V, Dekou V, Nicaud V, Chasse JF, London J, Stansbie D, Humphries SE, and Gudnason V; and Variation at the von willebrand factor (vWF) gene locus is associated with plasma vWF:Ag levels: identification of three novel single nucleotide polymorphisms in the vWF gene promoter. *Blood* (1999) **93**:4277-83, Keightley AM, Lam YM, Brady JN, Cameron CL, Lillicrap D).

**[0046]** According to another aspect of the present invention there is provided use of a statin in preparation of a medicament for treating a cardiovascular disease in a human detected as having an OATPC polymorphism at one or more of the following positions:

position 1561 of the OATPC gene as defined by the position in SEQ ID NO: 1 is not G;  
position 488 in OATPC polypeptide defined by position in SEQ ID NO: 2 is not Gly.

**[0047]** Preferably determination of the status of the human is clinically useful. Examples of clinical usefulness include deciding which statin drug or drugs to administer and/or in deciding on the effective amount of the statin drug or drugs. Statins already approved for use in humans include atorvastatin, cerivastatin, fluvastatin, pravastatin and simvastatin. The reader is referred to the following references for further information: Drugs and Therapy Perspectives (12<sup>th</sup> May 1997), **9**: 1-6; Chong (1997) Pharmacotherapy **17**: 1157-1177; Kellick (1997) Formulary **32**: 352; Kathawala (1991) Medicinal Research Reviews, **11**: 121-146; Jahng (1995) Drugs of the Future **20**: 387-404, and Current Opinion in Lipidology, (1997), **8**, 362 - 368. A preferred statin drug is compound 3a (S-4522) in Watanabe (1997) Bioorganic and Medicinal Chemistry **5**: 437-444; now called rosuvastatin, see Olsson (2001) American Journal of Cardiology, **87**, supplement 1, 33-36. The term "drug transportable by OATPC" means that transport by OATPC in humans is an important part of a drug exerting its pharmaceutical effect in man. For example, some statins have to be transported to the liver by OATPC to exert their lipid lowering effects.

**[0048]** According to another aspect of the present invention there is provided an allelic variant of human OATPC polypeptide comprising:

an arginine at position 488 of SEQ ID NO 2;  
or a fragment thereof comprising at least 10 amino acids provided that the fragment comprises at least one allelic variant.

**[0049]** Fragments of polypeptide are at least 10 amino acids, more preferably at least 15 amino acids, more preferably at least 20 amino acids.

**[0050]** The invention will now be illustrated but not limited by reference to the following Examples. All temperatures are in degrees Celsius.

**[0051]** In the Examples below, unless otherwise stated, the following methodology and materials have been applied.

**[0052]** AMPLITAQ™, available from Perkin-Elmer Cetus, is used as the source of thermostable DNA polymerase.

**[0053]** General molecular biology procedures can be followed from any of the methods described in "Molecular Cloning - A Laboratory Manual" Second Edition, Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory, 1989).

**[0054]** Electropherograms were obtained in a standard manner: data was collected by ABI377 data collection software and the wave form generated by ABI Prism sequencing analysis (2.1.2).

## Example 1

### **Identification of Polymorphisms**

#### **1. Methods**

##### DNA Preparation

**[0055]** DNA was prepared from frozen blood samples collected in EDTA following protocol I (Molecular Cloning: A Laboratory Manual, p392, Sambrook, Fritsch and Maniatis, 2<sup>nd</sup> Edition, Cold Spring Harbor Press, 1989) with the following modifications. The thawed blood was diluted in an equal volume of standard saline citrate instead of phosphate buffered saline to remove lysed red blood cells. Samples were extracted with phenol, then phenol/chloroform and then chloroform rather than with three phenol extractions. The DNA was dissolved in deionised water.

### Template Preparation

[0056] Templates were prepared by PCR using the oligonucleotide primers and annealing temperatures set out below. The extension temperature was 72° and denaturation temperature 94°. Generally 50 ng of genomic DNA was used in each reaction and subjected to 35 cycles of PCR. Where described below, the primary fragment was diluted 1/100 and two microlitres were used as template for amplification of secondary fragments. PCR was performed in two stages (primary fragment then secondary fragment) to ensure specific amplification of the desired target sequence.

Polymorphisms in OATPC: cDNA screening of 15 Liver samples				
Region	SNP	Position	Amino Acid Change	Allele frequencies
Exon 4	G/A	510	None	G=96.7% A=3.3%
Exon 5	C/T	670	None	C=50% T=50%
Exon 5	C/T	696	None	C=60% T=40%
Exon 9	C/G	1299	Phe400Leu	C=96.7% G=3.3%
Exon 9	G/A	1312	Va1405Ile	G=96.7% A=3.3%
Exon 9	G/A	1347	None	G=96.7% A=3.3%
Exon 10	G/C	1561	Gly488Arg	G=96.7% C=3.3%
Exon 14	A/C	2028	Leu643Phe	A=90% C=10%

[0057] OATP2 above refers to the clone sequenced by Hsiang et al (ref 1). Some comment on the numbering of exons in OATPC is required. This gene contains an exon (38 bp) upstream (5' UTR region) of the exon containing the ATG start site for translation. Therefore the exon numbering could vary depending whether this exon is counted as the first exon or not. In the literature, Konig (2000) JBC 275: 23161-68, have defined exon 1 as that containing the ATG start site and therefore we have adopted the same numbering in this application (but note that the priority document relating to the present application did vice versa; for example, exon 5 in this application is equivalent to exon 6 in the priority document).

PCR PRODUCTS		
Fragment	Forward Oligo	Reverse Oligo
443-999	443-466	979-999
874-1360	874-896	1337-1360
1255-1684	1255-1278	1663-1684
1559-2095	1559-1581	2073-2095

RFLP analysis			
Polymorphism	Position	RFLP Enzyme/PCR size	RFLP fragment size
G/A	510		

# EP 1 186 672 B1

(continued)

RFLP analysis			
Polymorphism	Position	RFLP Enzyme/PCR size	RLFP fragment size
C/T	670	BmR I/ 595bp	C=349bp, 246bp, T=595bp
C/T	696		
C/G	1299	Apo I/ 595bp	C=30bp, 60bp, 380 bp G=90bp, 380bp
G/A	1312	Bst 4Cl/ 595bp	G=77bp, 393bp A=470bp
G/A	1347		
G/C	1561	HpyCH4IV/ 595bp	G=470bp C=144bp, 326bp
A/C	2028	Ase I/ 595bp	A=89bp, 488bp C=577bp

Example 2

[0058]



<b>Further OATPC Polymorphisms</b>							
<b>SNPs in OATPC 3'UTR</b> (positions according to SEQ ID NO 1)							
<b>Exon</b>	<b>Nucleotide</b>	<b>SNP</b>				<b>Frequency</b>	<b>Frequency</b>
						<b>Caucasian</b>	<b>Japanese</b>
3' UTR	2327	Ins T				not screened	0.1
3' UTR	2342	T>C				not screened	0.4
<b>SNPs in OATPC promoter</b> (positions according to SEQ ID NO 3)							
	<b>Nucleotide</b>	<b>SNP</b>				<b>Frequency</b>	<b>Frequency</b>
						<b>Caucasian</b>	<b>Japanese</b>
	321	T>G				0.03	not screened
	1332	A>C				0.08	not screened
<b>SNPs in OATPC introns</b>							
<b>Intron</b>	<b>Nucleotide</b>	<b>SNP</b>	<b>Nucleotide</b>	<b>Sequence</b>			
	<b>position in</b>		<b>position</b>	<b>ID No</b>			
	<b>relation to exon</b>		<b>in sequence</b>				
1	IVS1+21	T>A	41	4			
2	IVS2+89	T>G	109	5			
2	IVS2+224	A>G	244	5			
3	IVS3+97	C>A	117	6			
3	IVS3+263	G>A	283	6			
4	IVS4+189	G>A	209	7			
4	IVS4+191	G>A	211	7			
4	IVS5-118	delCTTGTA	63	8			
6	IVS6+33	C>T	53	9			
9	IVS10-107	ins TTC	75	10			
11	IVS11+142	Ins T	162	11			
12	IVS13-97	G>C	84	12			

**OATPC intronic SNPs****Key**

[0059] 20bp of exon sequence shown in uppercase

Intron sequence in lowercase (200 to 300bp only)

SNP shown in uppercase (one allele only)

5 **Sequence ID No 4**  
**IVS1+21 T>A SNP at position 41 in this sequence**

GATACTGCAA TGGATTGAAG gtagaataag ttttatgttt Ttgagctaaa ataagtaaat 60  
 agggaaacttt aatgtataga aaagcaagtt gttaaaaaga acattatgtt tcaaattata 120  
 10 attttcaatt gaagcatata ttgaaatatt aacataatga ttcatacctt gattttaaac 180  
 agtcttttaa tctgattaag 200

15 **Sequence ID No 5**  
**IVS2+89 T>G SNP at position 109 in this sequence**  
**IVS2+224 A>G SNP at position 244 in this sequence**

TGACGGAAGC TTTGAAATTG gtaacatttta ttttctattt taataaccaa acttgcaaag 60  
 ttaaaaaata tatatgcttt acaccactgg ttatcaactg gggtaaatTt atctctcaca 120  
 20 ggcaatttgg caataactaa aaacatttgg gggtgtcata actgcacagg gggtgggggc 180  
 aatggaagtg ctactggtat ctaaaggtag aggtcagggg tactgctaaa tattctataa 240  
 tgcAcaaaga atgatgtaac tgaaaatggt gatagtggag atgttcagaa accctgattc 300

25 **Sequence ID No 6**  
**IVS3+97 C>A SNP at position 117 in this sequence**  
**IVS3+263 G>A SNP at position 283 in this sequence**

30 CCACATTTCT TCATGGGATA gtaagtgtta aaaaaaaaaa aaacctctgt gccactatca 60  
 gtaccttgta aattaggagt agaattttat tattatccct ttaaattaggc agttacCttt 120  
 tgagaagata cccactaagt gtgtacagaa atgaaatagt gtctatttgt ctacataatc 180  
 attttattta tcgtagcttt catatacttt gaaataacaa aaagactaaa ctgtagagtt 240  
 35 tcaaatgaaa taaataggct ttttatgaat ttttagtata acGtatatac tgtacgtcct 300

40 **Sequence ID No 7**  
**IVS4+189 G>A SNP at position 209 in this sequence**  
**IVS4+191 G>A SNP at position 211 in this sequence**

ACCTGAGATA GTGGGAAAAG gtaagaatta atattgacag taaaaagtct tctaaaatgt 60  
 atacatttaa ttacatctct aaaaattggt gtgatattca ttagcaaat ttaattaaga 120  
 atgaatagga aaaacatttg actcttacag acataattat agtgtaata tacacagttc 180  
 gcccat AAC aacacaggtt taaactacGc Gttttcactt ctatgcaaat tttgtccatc 240  
 45 tgaactggat gataaacctg ccggtaagaa tatctgacat tttctatatt tggattgaac 300

50 **Sequence ID No 8**  
**IVS5-118 delCTTGTA deletion of 6bp from position 63 to 68 incl.**

tagcagcata agaattggact aatacaccat attgtcaaag tttgcaaagt gaatataaat 60  
 taCTTGTAAct tgtaaattaa aaaaaataa gtagaataat taagagttaa caagtagtta 120  
 aatttgtaat agaaatgcta aaattaatgt ttaaaatgaa acactctctt atctacatag 180  
 GTTGTTTAAA GGAATCTGGG 200

55

Sequence ID No 9 IVS6+33 C>T                      SNP at position 53 in this sequence  
TATTGGATAT GTAGATCTAA gtaagtacaa ccagaacaag gtaccatgat aaCgtctttc 60  
taagcacaca tgcgaaaaac attttttcaa ataactgaat tcactctttc aatagtcctt 120  
5 tgcttaatat aattagaaag ttacaagtag gaaataaatg tattactaat cagaataaat 180  
ataaaatcca gctcctattt 200

Sequence ID No 10  
10 IVS10-107 Ins TTC                      SNP at position 75 in this sequence  
ttaaaaaaaaa ctttgccatt tcgtcatcat caaagcaaatt ttcttcatat aaagaaaaat 60  
tctttatcta cttt(TTC)ttttcc ctctttctct gctttcactt tacttcttcc ttctcctccc 120  
cttctttgtc tttttcttct ctctctctct ttttgatata tgtctatcat atatttccag 180  
AAATAATCCA GTGACATCTC 200

Sequence ID No 11  
15 IVS11+142 InsT                      InsT at position 162 in this sequence  
CATGTCATGC TGATTGTTAA gtaagtatga cttttaaaaa cattttcata tgcattgagac 60  
20 tataaacaca cctaattgata tgcataattt tacataatat actgggaatt caaattcata 120  
tttcatcaaa ttttaatttt ctgagaattc attttattaa aa(T)ttactatg aactctcaag 180  
gctgtaatta ataattttgc 200

Sequence ID No 12  
25 IVS13-97 G>C                      SNP at position 84 in this sequence  
tgatttgggt ctttgagatt tctaataatc ttattatttg ggtagatgca gaacaaaaata 60  
ataaacgaat cctccaaatt tttGaacttt tatttaataa aaatatatca atgtggaata 120  
30 tcatgcagtt acatttaaaa tatgttccct aaactgacat cttctcttct cctattacag 180  
GAGGAATTCT AGCTCCAATA 200

#### OATPC promoter region

35 [0060] Total length of the sequence = 1538bp  
1500bp of OATPC sequence directly upstream from the cDNA sequence  
Sequence in uppercase represents 38bp overlap with the cDNA sequence (SEQ ID NO 1) where this 38bp is 5'UTR  
sequence.  
Nucleotide positions in the promoter have been determined where the -1 position is the base (lowercase) directly  
40 upstream of the end of the cDNA sequence.

# EP 1 186 672 B1

```

atgctctttg acctctgaaa atattggaga attttacaac tggcaccttt agctcaggat 60
tataaagggt gttagttagt ttgtactggt ttatcttcat tgtatataat atatatatta 120
gtctccaaac atgttgatgt gttttcaatg aaatggatgt ctgaggagaa aaccattagc 180
5 ctgagaaaac ccaaactgta ttcccatgtt gaataaaaagg aagtccataa aaatgatgga 240
aaatgttctg cattcctggt atgatatcaa aatctggcag tacatgaaaa tttttcaaag 300
tgcttattta acaggcataa tctttgggtc cctgagccag aatctgctgg gtatgggact 360
ggattgctat tttgacaact cggcagtaga ttcttactca gcagagtatt tgggaagcctt 420
actctaatat tttggccttg ggtctacatt tctcagttct gcacagtcac tcttcccctc 480
tacactactc tttagtgtgt ctcatgattc caatactctc aataattaac caagaataga 540
10 actaatcaat cagataactg tggcacagac atcaaataca ttttgctgca accatatcaa 600
caaatgtccc atgaatgata aggggtaacc atattctcat atatgcatcc tcacattacc 660
acatatatat atgtgcatat gtgtatacag gtaaaagtgt gtatatatgt atacatgtat 720
gtttgtgtgt atatacatat atatatcttc acacttttct gaaatatata tatttatgtg 780
agagaagggt ctgtacttta tttcagaaga gagcttaatg tccaagggtat aattgagagt 840
ctaaaatggt tgagtatttg aattaatata acttcatctc tactcaagaa aacttttaac 900
tgagttaagc tcttcctttc tccacaagtc aagtcaataa aaggaaactg tgatattaat 960
15 aattctttcc tgttttgatg taaagaatct atcgcataaa gcagtcttaa ttttcatcat 1020
tcagaaaaat ggtcttgcag ttaattggga ctctcttatt ccaggtggtt tctccagtct 1080
ccatacatat cacgtagtaa ccatacttat gtaccaagca aagagggtat attttaattt 1140
ttaaatgcca atgtaacctg taggcatatt ttttatttgt cttaaattat ttctatttg 1200
gaagttttaa atacctggaa taatttattg tactcatatt tttaaagaaa aaaatcttat 1260
gccaccaact taattgaata aacaagtaaa agccattccc aaaagtaagg ttacttgtt 1320
20 aagattaaac aaaaataatg tgagaattct gagaaatata atctttaa attggaact 1380
ggagtgaact cttaaaacta actagggttt atatgtttga ctgagcaat gacataataa 1440
ggtggttaat catcactgga cttgttttca aaaagccaac tactttaaga ggaataaagg 1500
GTGGACTTGT TGCAGTTGCT GTAGGATTCT AAATCCAG 1538

```

## SEQUENCE LISTING

[0061]

```

30      <110> AstraZeneca AB

      <120> Chemical Compounds

35      <130> adeokun
           ambrose
           cresswell
           dudley

40      <140>
      <141>

      <160> 12

45      <170> PatentIn Ver. 2.1

           <210> 1
           <211> 2452
           <212> DNA
50      <213> Homo sapiens

           <400> 1

```

EP 1 186 672 B1

5  
 10  
 15  
 20  
 25  
 30  
 35  
 40  
 45  
 50  
 55

gtggacttgt	tgcagttgct	gtaggattct	aaatccaggt	gattgtttca	aactgagcat	60
caacaacaaa	aacatttgta	tgatatctat	atctcaatca	tggaccacaaa	tcaacatttg	120
aataaaacag	cagaggcaca	accttcagag	aataagaaaa	caagatactg	caatggattg	180
aagatgttct	tggcagctct	gtcactcagc	tttatgtgta	agacactagg	tgcaattatt	240
atgaaaagtt	ccatcattca	tatagaacgg	agatttgaga	tatcctcttc	tcttggtggg	300
tttattgacg	gaagctttga	aattggaaat	ttgcttgtga	ttgtatttgt	gagttacttt	360
ggatccaaac	tacatagacc	aaagttaatt	ggaatcggtt	gtttcattat	gggaattgga	420
gggtttttga	ctgctttgcc	acatttcttc	atgggatatt	acaggtattc	taaagaaact	480
aatatcaatt	catcagaaaa	ttcaacatcg	accttatcca	cttggtttaat	taatcaaat	540
ttatcactca	atagagcatc	acctgagata	gtgggaaaaag	gttggtttaa	ggaatctggg	600
tcatacatgt	ggatatatgt	gttcatgggt	aatatgcttc	gtggaatagg	ggagactccc	660
atagtaccac	tggggctttc	ttacattgat	gatttcgcta	aagaaggaca	ttcttctttg	720
tatttaggta	tattgaatgc	aatagcaatg	attggtccaa	tcattggcct	tacctggga	780
tctctgtttt	ctaaaatgta	cgtggatatt	ggatatgtag	atctaagcac	tatcaggata	840
actcctactg	attctcgatg	gggtggagct	tgggtggctta	atttccttgt	gtctggacta	900
ttctccatta	tttcttccat	accattcttt	ttcttgcccc	aaactccaaa	taaaccacaa	960
aaagaaagaa	aagcttcact	gtctttgcat	gtgctggaaa	caaatgatga	aaaggatcaa	1020
acagctaatt	tgaccaatca	aggaaaaaat	attaccaaaa	atgtgactgg	ttttttccag	1080
tcttttaaaa	gcacccctac	taatccctcg	tatgttatgt	ttgtgctttt	gacgttggtta	1140
caagtaagca	gctatattgg	tgcttttact	tatgtcttca	aatacgtaga	gcaacagtat	1200
ggtcagcctt	catctaaggc	taacatctta	ttgggagtca	taaccatacc	tatttttgca	1260
agtggaatgt	ttttaggagg	atatatcatt	aaaaaattca	aactgaacac	cgttggaatt	1320
gccaaattct	catgttttac	tgctgtgatg	tcattgtcct	tttacctatt	atattttttc	1380
atactctgtg	aaaacaaatc	agttgccgga	ctaaccatga	cctatgatgg	aaataatcca	1440
gtgacatctc	atagagatgt	accactttct	tattgcaact	cagactgcaa	ttgtgatgaa	1500
agtcaattgg	aaccagctctg	tggaaacaat	ggaataaact	acatctcacc	ctgtctagca	1560
ggttgcaaat	cttcaagtgg	caataaaaag	cctatagtgt	tttacaactg	cagttgtttg	1620
gaagtaactg	gtctccagaa	cagaaattac	tcagcccat	tgggtgaatg	ccaagagat	1680
gatgcttgta	caaggaaatt	ttactttttt	gttgcaatac	aagtcttgaa	tttatttttc	1740
tctgcacttg	gaggcacctc	acatgtcatg	ctgatgtgta	aaattgttca	acctgaattg	1800
aaatcacttg	cactgggttt	ccactcaatg	gttatacgag	cactaggagg	aattctagct	1860
ccaatatatt	ttggggctct	gattgataca	acgtgtataa	agtgtccac	caacaactgt	1920
ggcacacgtg	ggtcatgtag	gacatataat	tccacatcat	tttcaagggg	ctactggggc	1980
ttgtcttcaa	tgtaagagt	ctcatcactt	gttttatata	ttatattaat	ttatgccatg	2040
aagaaaaaat	atcaagagaa	agatatcaat	gcacagaaaa	atggaagtgt	catggatgaa	2100
gcaaaacttag	aatccttaaa	taaaaataaa	cattttgtcc	cttctgctgg	ggcagatagt	2160
gaaacacatt	gttaaggggga	gaaaaaaagc	cacttctgct	tctgtgtttc	caaacagcat	2220
tgcatgtgatt	cagtaagatg	ttatttttga	ggagtctctg	gtcctttcac	taagaatttc	2280
cacatctttt	atggtggaag	tataaataag	cctatgaact	tataataaaa	caaactgtag	2340
gtagaaaaaa	tgagagtact	cattgtacat	tatagctaca	tatttgtggg	taagggttaga	2400
ctatatgate	catacaaat	aaagtggag	acatgggttac	tgtgtaataa	aa	2452

<210> 2  
 <211> 691  
 <212> PRT  
 <213> Homo sapiens  
 <400> 2

EP 1 186 672 B1

	Met	Asp	Gln	Asn	Gln	His	Leu	Asn	Lys	Thr	Ala	Glu	Ala	Gln	Pro	Ser	
	1				5					10					15		
5	Glu	Asn	Lys	Lys	Thr	Arg	Tyr	Cys	Asn	Gly	Leu	Lys	Met	Phe	Leu	Ala	
				20				25						30			
	Ala	Leu	Ser	Leu	Ser	Phe	Ile	Ala	Lys	Thr	Leu	Gly	Ala	Ile	Ile	Met	
			35					40					45				
10	Lys	Ser	Ser	Ile	Ile	His	Ile	Glu	Arg	Arg	Phe	Glu	Ile	Ser	Ser	Ser	
		50				55						60					
	Leu	Val	Gly	Phe	Ile	Asp	Gly	Ser	Phe	Glu	Ile	Gly	Asn	Leu	Leu	Val	
	65				70					75						80	
	Ile	Val	Phe	Val	Ser	Tyr	Phe	Gly	Ser	Lys	Leu	His	Arg	Pro	Lys	Leu	
15					85					90					95		
	Ile	Gly	Ile	Gly	Cys	Phe	Ile	Met	Gly	Ile	Gly	Gly	Val	Leu	Thr	Ala	
				100					105					110			
	Leu	Pro	His	Phe	Phe	Met	Gly	Tyr	Tyr	Arg	Tyr	Ser	Lys	Glu	Thr	Asn	
20			115					120					125				
	Ile	Asn	Ser	Ser	Glu	Asn	Ser	Thr	Ser	Thr	Leu	Ser	Thr	Cys	Leu	Ile	
		130					135					140					
	Asn	Gln	Ile	Leu	Ser	Leu	Asn	Arg	Ala	Ser	Pro	Glu	Ile	Val	Gly	Lys	
25	145					150					155					160	
	Gly	Cys	Leu	Lys	Glu	Ser	Gly	Ser	Tyr	Met	Trp	Ile	Tyr	Val	Phe	Met	
				165						170					175		
	Gly	Asn	Met	Leu	Arg	Gly	Ile	Gly	Glu	Thr	Pro	Ile	Val	Pro	Leu	Gly	
30				180					185					190			
	Leu	Ser	Tyr	Ile	Asp	Asp	Phe	Ala	Lys	Glu	Gly	His	Ser	Ser	Leu	Tyr	
			195				200						205				
	Leu	Gly	Ile	Leu	Asn	Ala	Ile	Ala	Met	Ile	Gly	Pro	Ile	Ile	Gly	Phe	
35		210				215					220						
	Thr	Leu	Gly	Ser	Leu	Phe	Ser	Lys	Met	Tyr	Val	Asp	Ile	Gly	Tyr	Val	
	225				230						235					240	
	Asp	Leu	Ser	Thr	Ile	Arg	Ile	Thr	Pro	Thr	Asp	Ser	Arg	Trp	Val	Gly	
40				245						250					255		
	Ala	Trp	Trp	Leu	Asn	Phe	Leu	Val	Ser	Gly	Leu	Phe	Ser	Ile	Ile	Ser	
				260					265					270			
	Ser	Ile	Pro	Phe	Phe	Phe	Leu	Pro	Gln	Thr	Pro	Asn	Lys	Pro	Gln	Lys	
			275				280						285				
45	Glu	Arg	Lys	Ala	Ser	Leu	Ser	Leu	His	Val	Leu	Glu	Thr	Asn	Asp	Glu	
		290				295						300					
	Lys	Asp	Gln	Thr	Ala	Asn	Leu	Thr	Asn	Gln	Gly	Lys	Asn	Ile	Thr	Lys	
	305					310					315					320	
50	Asn	Val	Thr	Gly	Phe	Phe	Gln	Ser	Phe	Lys	Ser	Ile	Leu	Thr	Asn	Pro	
				325						330					335		

EP 1 186 672 B1

	Leu Tyr Val Met Phe Val Leu Leu Thr Leu Leu Gln Val Ser Ser Tyr	
	340 345 350	
5	Ile Gly Ala Phe Thr Tyr Val Phe Lys Tyr Val Glu Gln Gln Tyr Gly	
	355 360 365	
	Gln Pro Ser Ser Lys Ala Asn Ile Leu Leu Gly Val Ile Thr Ile Pro	
	370 375 380	
10	Ile Phe Ala Ser Gly Met Phe Leu Gly Gly Tyr Ile Ile Lys Lys Phe	
	385 390 395 400	
	Lys Leu Asn Thr Val Gly Ile Ala Lys Phe Ser Cys Phe Thr Ala Val	
	405 410 415	
15	Met Ser Leu Ser Phe Tyr Leu Leu Tyr Phe Phe Ile Leu Cys Glu Asn	
	420 425 430	
	Lys Ser Val Ala Gly Leu Thr Met Thr Tyr Asp Gly Asn Asn Pro Val	
	435 440 445	
20	Thr Ser His Arg Asp Val Pro Leu Ser Tyr Cys Asn Ser Asp Cys Asn	
	450 455 460	
	Cys Asp Glu Ser Gln Trp Glu Pro Val Cys Gly Asn Asn Gly Ile Thr	
	465 470 475 480	
25	Tyr Ile Ser Pro Cys Leu Ala Gly Cys Lys Ser Ser Ser Gly Asn Lys	
	485 490 495	
	Lys Pro Ile Val Phe Tyr Asn Cys Ser Cys Leu Glu Val Thr Gly Leu	
	500 505 510	
30	Gln Asn Arg Asn Tyr Ser Ala His Leu Gly Glu Cys Pro Arg Asp Asp	
	515 520 525	
	Ala Cys Thr Arg Lys Phe Tyr Phe Phe Val Ala Ile Gln Val Leu Asn	
	530 535 540	
35	Leu Phe Phe Ser Ala Leu Gly Gly Thr Ser His Val Met Leu Ile Val	
	545 550 555 560	
	Lys Ile Val Gln Pro Glu Leu Lys Ser Leu Ala Leu Gly Phe His Ser	
	565 570 575	
40	Met Val Ile Arg Ala Leu Gly Gly Ile Leu Ala Pro Ile Tyr Phe Gly	
	580 585 590	
	Ala Leu Ile Asp Thr Thr Cys Ile Lys Trp Ser Thr Asn Asn Cys Gly	
	595 600 605	
45	Thr Arg Gly Ser Cys Arg Thr Tyr Asn Ser Thr Ser Phe Ser Arg Val	
	610 615 620	
	Tyr Leu Gly Leu Ser Ser Met Leu Arg Val Ser Ser Leu Val Leu Tyr	
	625 630 635 640	
	Ile Ile Leu Ile Tyr Ala Met Lys Lys Lys Tyr Gln Glu Lys Asp Ile	
	645 650 655	
50	Asn Ala Ser Glu Asn Gly Ser Val Met Asp Glu Ala Asn Leu Glu Ser	
	660 665 670	
	Leu Asn Lys Asn Lys His Phe Val Pro Ser Ala Gly Ala Asp Ser Glu	
	675 680 685	
55	Thr His Cys	

690

<210> 3  
 <211> 1538  
 <212> DNA  
 <213> Homo sapiens

<400> 3

```

atgctctttg acctctgaaa. atattggaga attttacaac tggcaccttt agctcaggat 60
tataaagggt gttagttagt ttgtactgtt ttatcttcat tgtatataat atatatatta 120
gtctccaaac atgttgatgt gttttcaatg aaatggatgt ctgaggagaa aaccattagc 180
ctgagaaaaa ccaaactgta tccccattgt gaataaaaagg aagtccataa aaatgatgga 240
aaatggttctg cattcctgtt atgatatcaa aatctggcag tacatgaaaa tttttcaaag 300
tgcttattta acaggcataa tctttggctc cctgagccag aatctgctgg gtatgggact 360
ggattgctat ttgacaact cgcagtaga ttcttactca gcagagtatt tggaaacctt 420
actctaatat tttggccttg ggtctacatt tctcagttct gcacagtcac tcttcccttc 480
tacctacttc tttagtttgt ctcatgattc caataacttc aataattaac caagaataga 540
actaatcaat cagataactg tggcacagac atcaaataca ttttgctgca accatatcaa 600
caaatgtccc atgaatgata aggggtaacc atatttctcat atatgcatcc tcacattacc 660
acatatatat atgtgcata gtgtatacag gtaaaagtg gtatatatgt atacatgtat 720
gtttgtgtgt atatacatc atatatcttc acacttttct gaaatatata tatttatgtg 780
agagaagggc ctgtacttta tttcagaaga gagcttaatg tccaagggtat aattgagagt 840
ctaaaatggt tgagttattg aattaattaa acttcatctc tactcaagaa aacttttaac 900
tgagttaagc tcttccttc tccacaagtc aagtcataa aaggaaactg tgatattaat 960
aattctttcc tgttttgatg taaagaatct atcgcataaa gcagtcctta ttttcatcat 1020
tcagaaaaat ggtcttgtag ttaattggga ctctcttatt ccagggtgga tctccagctc 1080
ccatacatac caggttagaa ccatacttat gtaccaagca aagaggggtat attttaattt 1140
ttaaatgcca atgtaacctg taggcataat ttttatttgt cttaaattat ttctatttg 1200
gaagttttta atacctggaa taattttatt tactcatatt tttaaagaaa aaaatcttat 1260
gccaccaact taattgaata aacaagtaaa agccattccc aaaagtaagg ttactttgtt 1320
aagattaaca aaaaataatg tgagaattct gagaaatata atctttaaat attggcaact 1380
ggagtgaact cttaaaacta actaggtttt atatgtttga ctagagcaat gacataataa 1440
gggtggttaat catcactgga ctgtgtttca aaaagccaac tactttaaga ggaataaagg 1500
gtggacttgt tgacgttgct gtaggattct aaatccag 1538

```

<210> 4  
 <211> 200  
 <212> DNA  
 <213> Homo sapiens

<400> 4

```

gatactgcaa tggattgaag gtagaataag ttttatgttt ttgagctaaa ataagtaaat 60
agggaaacttt aatgtataga aaagcaagtt gttaaaaaga acattatgtt tcaaattata 120
attttcaatt gaagcatata ttgaaatatt aacataatga ttcatacctt gattttaaacc 180
agtcttttaa tctgattaag

```

<210> 5  
 <211> 300  
 <212> DNA  
 <213> Homo sapiens

<400> 5



# EP 1 186 672 B1

```

5      tgacggaagc tttgaaattg gtaacattta ttttctatct taataaccaaa acttgcaaag 60
      ttaaaaaata tatatgcttt acaccactgg ttatcaactg gggtaaattt atctctcaca 120
      ggcaatttgg caataactaa aaacatttgg ggttgtcata actgcacagg ggttgggggc 180
      aatggaagtg ctactgggat ctaaaggtag aggtcagggg tactgctaaa tattctataa 240
      tgcacaaaga atgatgtaac tgaaaatgtt gatagtgagg atgttcagaa accctgattc 300

```

```

<210> 6
<211> 300
10    <212> DNA
      <213> Homo sapiens

```

```

<400> 6

```

```

15      ccacatttct tcatgggata gtaagtgtta aaaaaaaaaa aaacctctgt gccactatca 60
      gtaccttgta aattaggagt agaattttat tattatccct ttaaataaggc agttaccttt 120
      tgagaagata cccactaagt gtgtacagaa atgaaatagt gtctatttgt ctacataatc 180
      attttattta tcgtagcttt catatacttt gaaataacaa aaagactaaa ctgtagagtt 240
      tcaaatgaaa taaataggct ttttatgaat ttttagtata acgtatatac tgtacgtctt 300

```

```

25    <210> 7
      <211> 300
      <212> DNA
      <213> Homo sapiens

```

```

<400> 7

```

```

30      acctgagata gtgggaaaaag gtaagaatta atattgacag taaaaagtct tctaaaaatgt 60
      atacatttaa ttacatctct aaaaattgtt gtgatattca tttagcaaaat ttaattaaga 120
      atgaatagga aaaacatttg actcttacag acataattat agtggttaata tacacagttc 180
      gccattaac aacacaggtt taaactacgc gttttcactt ctatgcaaat tttgtccatc 240
      tgaactggat gataaacctg ccgtaagaa tatctgacat tttctatatt tggattgaac 300

```

```

35    <210> 8
      <211> 200
      <212> DNA
      <213> Homo sapiens

```

```

40    <400> 8

```

```

45      tagcagcata agaatggact aatacaccat attgtcaaag tttgcaaagt gaatataaat 60
      tacttgtact tgtaaattaa aaaaaataaa gtagaataat taagagttta caagtagtta 120
      aatttgtaat agaaatgcta aaattaatgt ttaaaatgaa acactctctt atctacatag 180
      gttgtttaa ggaatctggg                200

```

```

50    <210> 9
      <211> 200
      <212> DNA
      <213> Homo sapiens

```

```

<400> 9

```

```

55      tattggatat gtagatctaa gtaagtacaa ccagaacaag gtaccatgat aacgtctttc 60
      taagcacaca tgcgaaaaac attttttcaa ataactgaat tcactctttc aatagtcctt 120
      tgcttaatat aattagaaag ttacaagtag gaaataaatg tattactaat cagaataaat 180
      ataaaatcca gtccttattt                200

```

<210> 10  
 <211> 203  
 <212> DNA  
 <213> Homo sapiens

5

<400> 10

```

          ttaaaaaaaaa ctttgccatt tcgtcatcat caaagcaaatt ttcttcatat aaagaaaaat 60
          tctttatcta ctttttcttt tccctctttc tctgctttca ctttacttct cccttctcct 120
          ccccttcttt gtctttttct tctctctctc tctttttgat atatgtctat catatatttc 180
          cagaaataat ccagtacat ctc                                     203
  
```

10

<210> 11  
 <211> 201  
 <212> DNA  
 <213> Homo sapiens

15

<400> 11

20

```

          catgtcatgc tgattgttaa gtaagtatga cttttaaaaa cattttcata tgcattgagac 60
          tataaacaca cctaattgata tgcataatttt tacataatat actgggaatt caaattcata 120
          ; tttcatcaaa ttttaatttt ctgagaattc attttattaa aatttactat gaacttctcaa 180
          ggctgtaatt aataattttg c                                     201
  
```

25

<210> 12  
 <211> 200  
 <212> DNA  
 <213> Homo sapiens

30

<400> 12

35

```

          tgatttgggt ctttgagatt tctaataatc tttattattg ggtagatgca gaacaaaata 60
          ataaacgaat cctccaaatt tttgaacttt tatttaatca aaatatatca atgtggaata 120
          tcatgcagtt acatttaaaa tatgttcctt aaactgacat cttctcttct cctattacag 180
          gaggaattct agtccaata,                                     200
  
```

40

## Claims

45

1. A method for the detection of a polymorphism in OATPC in a human, which method comprises determining the sequence of the human at:

50

position 1561 of the OATPC gene as defined by the position in SEQ ID NO: 1 is not G; or  
 position 488 in OATPC polypeptide defined by position in SEQ ID NO: 2 is not Gly.

2. Use of a method as defined in claim 1 to assess the pharmacogenetics of a drug transportable by OATPC.
3. A polynucleotide comprising at least 20 bases of the human OATPC gene and comprising the allelic variant:

55

Region	variant	Position in SEQ ID NO	SEQ ID NO
Exon 10	C	1561	1

4. An ARMS allele specific primer or an allele-specific oligonucleotide probe selected from one of the following:
  - an ARMS allele specific primer; or
  - an allele-specific oligonucleotide probe,
  - capable of detecting a OATPC gene polymorphism as defined in claim 1.
5. Use of an OATPC polymorphism as defined in claim 1 as a genetic marker in a linkage study.
6. Use of a statin in preparation of a medicament for treating a cardiovascular disease in a human detected as having an OATPC polymorphism at one or more of the following positions:
  - position 1561 of the OATPC gene as defined by the position in SEQ ID NO: 1 is not G;
  - position 488 in OATPC polypeptide defined by position in SEQ ID NO: 2 is not Gly.
7. A use according to claim 6 in which the drug is rosuvastatin.
8. An allelic variant of human OATPC polypeptide comprising;
  - an arginine at position 488 of SEQ ID NO 2
  - or a fragment thereof comprising at least 10 amino acids provided that the fragment comprises the allelic variant.

#### Patentansprüche

1. Verfahren zum Nachweis eines Polymorphismus in OATPC in einem Menschen, wobei man in dem Verfahren bestimmt, daß die Sequenz des Menschen in:
  - Position 1551 des OATPC-Gens, wie durch die Position in SEQ ID NO: 1 definiert, nicht G ist; oder
  - Position 488 im OATPC-Polypeptid, definiert durch die Positionen in SEQ ID NO: 2, nicht Gly ist.
2. Verwendung eines Verfahrens gemäß Anspruch 1 zur Beurteilung der Pharmacogenetik eines durch OATPC transportierbaren Arzneistoffs.
3. Polynukleotid, umfassend wenigstens 20 Basen des menschlichen OATPC-Gens und umfassend die Allelvariante:

Region	Variante	Position in SEQ ID NO:	SEQ ID NO:
Exon 10	C	1561	1

4. ARMS-Allel-spezifischer Primer oder Allelspezifische Oligonukleotidsonde, ausgewählt aus einem der folgenden Moleküle:
  - einem ARMS-Allel-spezifischen Primer; oder
  - einer Allel-spezifischen Oligonukleotidsonde,
  - womit ein OATPC-Gen-Polymorphismus gemäß Anspruch 1 nachgewiesen werden kann.
5. Verwendung eines OATPC-Polymorphismus gemäß Anspruch 1 als genetischer Marker in einer Verknüpfungsuntersuchung.
6. Verwendung eines Statins bei der Herstellung eines Arzneimittels zur Behandlung einer Herz-Kreislauf-erkrankung in einem Menschen, bei dem ein OATPC-Polymorphismus an einer oder mehreren der folgenden Positionen nachgewiesen wurde:
  - Position 1561 des OATPC-Gens, wie durch die Positionen in SEQ ID NO: 1 definiert, ist nicht G;
  - Position 488 im OATPC-Polypeptid, definiert durch die Positionen in SEQ ID NO: 2, ist nicht Gly.
7. Verwendung nach Anspruch 6, wobei es sich bei dem Arzneistoff um Rosuvastatin handelt.
8. Allelvariante des menschlichen OATPC-Polypeptids, umfassend:

ein Arginin in Position 488 der SEQ ID NO: 2  
oder ein Fragment davon, umfassend wenigstens 10 Aminosäuren, vorausgesetzt, daß das Fragment die Allelvariante umfaßt.

5

## Revendications

1. Procédé pour la détection d'un polymorphisme dans un OATPC d'un être humain, lequel procédé comprend l'étape consistant à déterminer la séquence de l'humain au niveau de :

10

la position 1561 du gène d'OATPC, comme cela est défini par la position dans la SEQ ID n° : 1, qui n'est pas un G ; ou  
la position 488 dans un polypeptide d'OATPC, définie par la position dans la SEQ ID n° : 2, qui n'est pas une Gly.

2. Utilisation d'un procédé, comme cela est défini à la revendication 1, pour évaluer les aspects pharmacogénétiques d'une substance médicamenteuse pouvant être transportée par un OATPC.

3. Polynucléotide comprenant au moins 20 bases du gène d'OATPC humain et comprenant le variant allélique :

20

Région	Variant	Position dans la SEQ ID n° :	SEQ ID n° :
Exon 10	C	1561	1

4. Amorce spécifique d'un allèle ARMS ou sonde oligonucléotidique spécifique d'un allèle choisie parmi l'un des éléments suivants :

une amorce spécifique d'un allèle ARMS ; ou  
une sonde oligonucléotidique spécifique d'un allèle,  
capable de détecter un polymorphisme dans un gène d'OATPC, comme cela est défini à la revendication 1.

30

5. Utilisation d'un polymorphisme d'OATPC, comme cela est défini à la revendication 1, en tant que marqueur génétique dans une étude de liaison.

6. Utilisation d'une statine dans la préparation d'un médicament destiné à traiter une affection cardiovasculaire chez un être humain, chez qui on a détecté un polymorphisme d'OATCP au niveau d'une ou plusieurs positions parmi les suivantes :

35

la position 1561 du gène d'OATPC, comme cela est défini par la position dans la SEQ ID n° : 1, n'est pas un G ; ou  
la position 488 dans un polypeptide d'OATPC, définie par la position dans la SEQ ID n° : 2, n'est pas une Gly.

40

7. Utilisation selon la revendication 6, dans laquelle la substance médicamenteuse est la rosuvastatine.

8. Variant allélique d'un polypeptide d'OATPC humain comprenant :

45

une arginine à la position 488 de la SEQ ID n° : 2 ou un fragment de celle-ci comprenant au moins 10 acides aminés, à condition que le fragment comprenne le variant allélique.

50

55